

Robustness of coherence: An operational and observable measure of quantum coherence

Carmine Napoli,^{1,2} Thomas R. Bromley,² Marco Cianciaruso,^{1,2} Marco Piani,³ Nathaniel Johnston,⁴ and Gerardo Adesso²

¹*Dipartimento di Fisica “E. R. Caianiello”, Università degli Studi di Salerno, Via Giovanni Paolo II, I-84084 Fisciano (SA), Italy; and INFN Sezione di Napoli, Gruppo Collegato di Salerno, Italy*

²*School of Mathematical Sciences, The University of Nottingham, University Park, Nottingham NG7 2RD, UK*

³*SUPA and Department of Physics, University of Strathclyde, Glasgow G4 0NG, UK*

⁴*Department of Mathematics and Computer Science, Mount Allison University, Sackville, New Brunswick E4L 1E2, Canada*

(Dated: January 11, 2016)

Quantifying coherence is an essential endeavour for both quantum foundations and quantum technologies. Here the robustness of coherence is defined and proven a full monotone in the context of the recently introduced resource theories of quantum coherence. The measure is shown to be observable, as it can be recast as the expectation value of a coherence witness operator for any quantum state. The robustness of coherence is evaluated analytically on relevant classes of states, and an efficient semidefinite program that computes it on general states is given. An operational interpretation is finally provided: the robustness of coherence quantifies the advantage enabled by a quantum state in a phase-discrimination task.

PACS numbers: 03.65.Ud, 03.67.Bg, 03.67.Ac, 03.65.Ta

Nearly one century old, quantum mechanics is now livelier than ever. Fundamental experiments have just been able to demonstrate, beyond any major loophole, that quantum correlations are incompatible with a local realistic interpretation [1–3]. On the other hand, the realization that quantum properties can be harnessed for practical applications [4] is presently fuelling a heated international race [5] to develop and deploy quantum technologies [6]. This is no coincidence: the improved study and test of fundamental quantum properties and our increased ability to exploit them go hand in hand.

The most essential feature signifying quantumness in a single system and underpinning all forms of quantum correlations in composite systems [7–9] is *quantum coherence*, namely the possibility of creating superpositions of a set of orthogonal states. Revealing quantum coherence in the state of a natural complex or man-made device earmarks its behaviour as genuinely nonclassical [10, 11]. Its degree of coherence often quantifies the ability of such an object to be an effective medium for quantum-enhanced applications [12, 13], ranging from cryptography [14] to metrology [15] and thermodynamics [16, 17]. Thus, it has become imperative to accomplish a rigorous operational characterization of quantum coherence.

Recently, various approaches have been put forward to develop a resource theory of coherence [12, 13, 18–25]. A resource theory is defined by the notions of free states (i.e. those not containing the resource, and assumed available at no cost) and free operations (i.e. those one is restricted to, and that cannot transform free states into resource states) [26, 27]. Fixing a reference basis, which we can identify as the computational basis $|j\rangle$ without loss of generality (with $j = 0, \dots, d-1$ for a d -dimensional system), the set \mathcal{S} of free states in all such resource theoretic approaches to coherence is given by incoherent states diagonal in the reference basis, $\delta = \sum_{j=0}^{d-1} \delta_j |j\rangle\langle j|$. Notice that \mathcal{S} is a convex zero-measure subset of the set of all states. Any state ρ can be reduced to an incoherent one by a full dephasing operation Δ , which maps ρ into its diagonal part $\Delta(\rho) = \sum_{j=0}^{d-1} |j\rangle\langle j| \rho |j\rangle\langle j|$ in the reference basis.

However, different authors have considered different options in analyzing limitations on the processing of coherence. We mention the following alternative choices of free operations, in order of inclusion: incoherence preserving operations [12] \supset incoherent operations [13] \supset strictly incoherent operations [28] \supset translationally invariant operations [20] \supset genuinely incoherent operations [25]. By incoherence preserving operations we refer to the maximal set of quantum channels Λ^M which map incoherent states into incoherent states [12], i.e. $\Lambda^M(\delta) \in \mathcal{S}$ for any $\delta \in \mathcal{S}$. Incoherent operations are instead defined as those quantum channels Λ^I which admit one operator sum decomposition $\Lambda^I(\rho) = \sum_l K_l \rho K_l^\dagger$ where the Kraus operators $\{K_l\}$ are all incoherence preserving subchannels [13]. Strictly incoherent operations Λ^S are a subset of incoherent operations admitting a set of incoherence preserving Kraus operators $\{K_l\}$ which further obey $\langle j|K_l \rho K_l^\dagger|j\rangle = \langle j|K_l \Delta(\rho) K_l^\dagger|j\rangle \forall j, l$, meaning that they can neither create nor make use of coherence [19, 28]. Even more restrictively, genuinely incoherent operations Λ^G are those which leave every incoherent state invariant, $\Lambda^G(\delta) = \delta$ [25]; for these channels (also known as generalized incoherent measurements [28]), according to all possible operator-sum decompositions, the Kraus operators are all incoherence preserving. Intermediate between the last two sets are translationally invariant operations, introduced within the complementary resource theory of asymmetry [18, 20]: specialized to coherence (i.e., asymmetry with respect to time translations generated by a Hamiltonian H diagonal in the reference basis $\{|j\rangle\}$), translationally invariant operations Λ^T are defined by the condition $e^{-iHt} \Lambda^T(\rho) e^{iHt} = \Lambda^T(e^{-iHt} \rho e^{iHt})$ for any ρ and any real t .

Several quantities have been proposed accordingly as candidate measures of quantum coherence, respecting physical requirements of monotonicity under (some of) the sets of operations introduced above [12, 13, 18–25, 29–31]. A canonical measure which complies with all the aforementioned resource theories is the relative entropy of coherence [12, 13, 18], which for a state ρ takes the simple form $\mathcal{C}_S(\rho) =$

$S(\Delta(\rho)) - S(\rho)$, where $S(\rho) = -\text{Tr}[\rho \log_2 \rho]$ is the von Neumann entropy. This measure can be interpreted as the optimal rate of maximally coherent states that can be distilled by incoherent operations Λ^I in the asymptotic limit of many copies of ρ [23]; however its experimental determination requires full state tomography, which can be unfeasible for large systems. More accessible measures of practical relevance for quantum metrology [15], such as the Wigner-Yanase skew information and the quantum Fisher information [29, 30], are monotone under translationally invariant operations but not under the larger set of incoherent operations [20], which may put into question their broader status as coherence quantifiers. In general, despite the pivotal role played by coherence in quantum information processing and the recent progress on its characterization, there remains a shortage of mathematically rigorous and physically intuitive bona fide measures of coherence endowed with direct operational interpretations.

In this Letter we fill this gap by introducing the *robustness of coherence*. As the name suggests, it quantifies the minimal mixing required to destroy all the coherence in a quantum state — an already operational definition, inspired by similar concepts previously introduced for entanglement, steering-type correlations, non-locality and even post-quantum correlations [32–35], and in general meaningful in any resource theory [27]. We prove that such a measure is a full monotone in all possible resource theories of coherence. The measure is furthermore computable (exactly in relevant cases, and numerically in general via a simple semidefinite program [36]) and observable: it can be recast as the expectation value of a witness operator for any quantum state. This makes it very appealing for experimental investigations, e.g. in condensed matter and biological contexts [10, 11, 37]. We then show that the measure admits a direct operational interpretation: it quantifies the advantage enabled by a quantum state, compared to any incoherent state, in a phase-discrimination task.

Let us remark that the robustness of coherence, with its outlined applications, can be generalized to the notion of robustness of asymmetry for arbitrary groups; these developments are explored in a companion paper [38]. The latter also contains detailed proofs for some technical results of this Letter.

Let $\mathcal{D}(\mathbb{C}^d)$ be the convex set of density operators acting on a d -dimensional Hilbert space, and let $\mathcal{I} \subset \mathcal{D}(\mathbb{C}^d)$ be the subset of incoherent states. We define the robustness of coherence (RoC) of a quantum state $\rho \in \mathcal{D}(\mathbb{C}^d)$ as

$$\mathcal{C}_{\mathcal{R}}(\rho) = \min_{\tau \in \mathcal{D}(\mathbb{C}^d)} \left\{ s \geq 0 \mid \frac{\rho + s\tau}{1+s} =: \delta \in \mathcal{I} \right\}, \quad (1)$$

that is, the minimum weight of another state τ such that its convex mixture with ρ yields an incoherent state δ . The concept is illustrated in Fig. 1 for a qubit ($d = 2$). If we denote by τ^* and δ^* the states achieving the minimum in (1), then

$$\rho = (1 + \mathcal{C}_{\mathcal{R}}(\rho))\delta^* - \mathcal{C}_{\mathcal{R}}(\rho)\tau^*, \quad (2)$$

is said to realize an optimal pseudomixture for ρ . Notice that it is necessary in Eq. (1) to let τ be an arbitrary state: if one




Figure 1. (Color online) Robustness of coherence $\mathcal{C}_{\mathcal{R}}(\rho)$ for a single qubit state $\rho = \frac{1}{2}(\mathbb{1} + \vec{r} \cdot \vec{\sigma})$, where \vec{r} is the Bloch vector and $\vec{\sigma}$ is the vector of Pauli matrices. Incoherent states span the thick vertical r_3 axis. The optimization in Eq. (1) is fulfilled by an equatorial pure state τ^* as depicted, resulting in $\mathcal{C}_{\mathcal{R}}(\rho) = (r_1^2 + r_2^2)^{\frac{1}{2}} = 2|\rho_{01}|$.

restricted τ to be incoherent, then the minimum s would diverge for any state ρ with nonzero coherence, henceforth resulting totally uninformative. This contrasts with the case of entanglement, for which the original robustness was defined in terms of pseudomixtures with separable states [32], and only later extended to pseudomixtures with arbitrary states [33].

Let us now prove that the RoC is a bona fide measure of coherence. First of all, it is seen by definition that

$$\mathcal{C}_{\mathcal{R}}(\rho) \geq 0 \quad \text{and} \quad \mathcal{C}_{\mathcal{R}}(\rho) = 0 \iff \rho \in \mathcal{I}. \quad (3)$$

Second, the RoC is convex, which, albeit not a fundamental request, is a desirable property for a coherence quantifier [13]. The proof follows closely the one for the robustness of entanglement [32]. Let ρ_1 and ρ_2 be two states, and consider for each the optimal pseudomixture $\rho_k = (1 + \mathcal{C}_{\mathcal{R}}(\rho_k))\delta_k^* - \mathcal{C}_{\mathcal{R}}(\rho_k)\tau_k^*$, with $k = 1, 2$. Take now the convex combination $\rho = p\rho_1 + (1-p)\rho_2$ with $0 \leq p \leq 1$, and notice that a pseudomixture $\rho = (1+s)\delta - s\tau$ can be written, with $\delta = [p(1 + \mathcal{C}_{\mathcal{R}}(\rho_1))\delta_1^* + (1-p)(1 + \mathcal{C}_{\mathcal{R}}(\rho_2))\delta_2^*]/(1+s) \in \mathcal{I}$, $\tau = [p\mathcal{C}_{\mathcal{R}}(\rho_1)\tau_1^* + (1-p)\mathcal{C}_{\mathcal{R}}(\rho_2)\tau_2^*]/s$, and $s = p\mathcal{C}_{\mathcal{R}}(\rho_1) + (1-p)\mathcal{C}_{\mathcal{R}}(\rho_2)$. By definition, $\mathcal{C}_{\mathcal{R}}(\rho) \leq s$, which proves convexity,

$$\mathcal{C}_{\mathcal{R}}(p\rho_1 + (1-p)\rho_2) \leq p\mathcal{C}_{\mathcal{R}}(\rho_1) + (1-p)\mathcal{C}_{\mathcal{R}}(\rho_2). \quad (4)$$

Third, and most importantly, the RoC is nonincreasing under all the aforementioned sets of operations used in resource theories of coherence. We prove in fact a general form of monotonicity under incoherence preserving (sub)channels. Let $\{\Gamma_l\}_{l=1}^m$ denote an instrument, i.e., a set of m (sub)channels (completely positive maps whose sum $\sum_{l=1}^m \Gamma_l(\rho) =: \Lambda(\rho)$ defines the action of a trace preserving channel Λ on ρ), which map any incoherent state $\delta \in \mathcal{I}$ into another (un)normalized incoherent state $\Gamma_l(\delta)$. For any ρ , we have then

$$\mathcal{C}_{\mathcal{R}}(\rho) \geq \sum_{l=1}^m \text{Tr}[\Gamma_l(\rho)] \mathcal{C}_{\mathcal{R}}\left(\frac{\Gamma_l(\rho)}{\text{Tr}[\Gamma_l(\rho)]}\right). \quad (5)$$

The proof can be easily sketched (see [38] for more details). Take the optimal pseudomixture for ρ given by Eq. (2)

and apply the (sub)channel Γ_l to both sides, $\Gamma_l(\rho) = (1 + \mathcal{C}_{\mathcal{R}}(\rho))\Gamma_l(\delta^*) - \mathcal{C}_{\mathcal{R}}(\rho)\Gamma_l(\tau^*)$. Since $\Gamma_l(\delta^*)/\text{Tr}[\Gamma_l(\delta^*)]$ is still incoherent, it is immediate to realize that the definition (1) of RoC implies that $\mathcal{C}_{\mathcal{R}}\left(\frac{\Gamma_l(\rho)}{\text{Tr}[\Gamma_l(\rho)]}\right) \leq \mathcal{C}_{\mathcal{R}}(\rho)\frac{\text{Tr}[\Gamma_l(\tau^*)]}{\text{Tr}[\Gamma_l(\rho)]}$. We have then $\sum_l \text{Tr}[\Gamma_l(\rho)]\mathcal{C}_{\mathcal{R}}\left(\frac{\Gamma_l(\rho)}{\text{Tr}[\Gamma_l(\rho)]}\right) \leq \sum_l \text{Tr}[\Gamma_l(\rho)]\mathcal{C}_{\mathcal{R}}(\rho)\frac{\text{Tr}[\Gamma_l(\tau^*)]}{\text{Tr}[\Gamma_l(\rho)]} = \mathcal{C}_{\mathcal{R}}(\rho)\sum_l \text{Tr}[\Gamma_l(\tau^*)] = \mathcal{C}_{\mathcal{R}}(\rho)$, concluding the proof. In the case $m = 1$, Eq. (5) proves that the RoC cannot increase, on average, under the maximal set of incoherence preserving operations $\{\Lambda^M\}$ [12]. For $m \geq 1$, if one identifies each Γ_l with a Kraus operator K_l (obeying $\sum_{l=0}^{m-1} K_l^\dagger K_l = \mathbb{1}$), then Eq. (5) proves instead monotonicity of the RoC under selective incoherent operations $\{\Lambda^I\}$, flagged as property C2b in [13], which is typically a rather demanding requirement in the framework of resource theories. Overall, Eq. (5) establishes therefore that the RoC is a *full monotone* with respect to *all* currently proposed formulations of the theory of quantum coherence.

We now show that the RoC has also desirable properties of computability and accessibility. Inspired by entanglement witnesses [7, 39], which are very useful tools to detect inseparability in laboratory [40], we introduce the notion of *coherence witness*. A Hermitian operator W satisfies $\Delta(W) \geq 0$ if and only if $\text{Tr}[\delta W] = \text{Tr}[\delta\Delta(W)] \geq 0$ for all incoherent states $\delta \in \mathcal{I}$; we call any such operator W a coherence witness, because finding $\text{Tr}[\rho W] < 0$ reveals the presence of coherence in the state ρ [41]. It turns out that the expectation value of any witness W , obeying the further constraint $W \leq \mathbb{1}$, provides a quantitative lower bound to the RoC [38], that is,

$$\max\{0, -\text{Tr}[\rho W]\} \leq \mathcal{C}_{\mathcal{R}}(\rho), \quad \forall W \text{ such that} \quad (6)$$

$$\Delta(W) \geq 0 \text{ and } W \leq \mathbb{1}. \quad (7)$$

Importantly, given any state ρ , there always exists an optimal witness W^* , characterized in particular by $\Delta(W^*) = 0$, which saturates inequality (6). In other words, the RoC is an *observable* quantity, given by the expectation value of a suitable (state-dependent) witness operator for any quantum state ρ . Finding such an optimal witness, hence determining $\mathcal{C}_{\mathcal{R}}(\rho)$ as defined in (1), can be then recast [38] as a simple and efficient semidefinite program [42], similar to the convex optimization one for the robustness of entanglement [43]:

$$\text{maximize } -\text{Tr}[W\rho] \quad \text{subject to Eq. (7)}. \quad (8)$$

For the convenience of the reader, we release MATLAB [44] code that makes use of the free CVX package [45, 46] to evaluate the RoC, as a Supplemental Material [36].

The above observations reveal that one can readily estimate the RoC from below in laboratory, by measuring any operator W obeying the constraints in (7), without the need for a full tomographical reconstruction of the density matrix ρ . This may be particularly valuable for investigations of coherence effects within biological domains, e.g. energy transport phenomena in light-harvesting systems [10, 11, 37, 47]. Notice however that, given a state ρ , the lower bound of Eq. (6) can vanish for non-optimized choices of W . In particular, one typically needs some knowledge on the form of ρ to determine the optimal witness W^* ; this issue is reminiscent of the case

of entanglement detection [40]. Nonetheless it is worth noticing that Eqs. (6) and (7) imply that, for any set of observables $\{O_i\}$, $i = 1, \dots, k$, measured in the lab with expectation values $o_i = \text{Tr}[O_i\rho]$, and not necessarily tailored to the measurement of RoC, one can consider coherence witnesses of the form $W = \sum_{i=1}^k c_i O_i + m\mathbb{1}$, for $c_1, \dots, c_k, m \in \mathbb{R}$, and obtain a lower bound to the RoC by the SDP [38] (code available [36])

$$\text{maximize } -(\sum_{i=1}^k c_i o_i + m)$$

$$\text{subject to } \Delta(\sum_{i=1}^k c_i O_i + m\mathbb{1}) \geq 0, \quad \sum_{i=1}^k c_i O_i + m\mathbb{1} \leq \mathbb{1}.$$

One can even make potentially better use of available experimental data, by exactly estimating the minimal RoC compatible with the data; this can also be cast as an SDP [36, 38].

Accessible *faithful* lower bounds to the RoC can be given too, noting that $W_2 = (\Delta(\rho) - \rho)/\|\Delta(\rho)\|_\infty$ obeys (7), so that

$$\mathcal{C}_{\mathcal{R}}(\rho) \geq \frac{\|\rho - \Delta(\rho)\|_2^2}{\|\Delta(\rho)\|_\infty} \geq \frac{\|\rho - \Delta(\rho)\|_2^2}{\|\Delta(\rho)\|_2} \geq \|\rho - \Delta(\rho)\|_2, \quad (9)$$

since $\text{Tr}[(\Delta(\rho) - \rho)\rho] = \text{Tr}[\Delta(\rho)^2] - \text{Tr}[\rho^2] = \|\rho - \Delta(\rho)\|_2^2$. Here, $\|\cdot\|_2$ is the 2-norm, and $\|\cdot\|_\infty$ is the operator norm. The quantity on the rightmost-hand side of (9) is: (i) nonzero on all but incoherent states; (ii) itself a monotone under genuinely incoherent operations Λ^G [25], but not under the larger sets of incoherent operations [13]; (iii) accessible via the measurement of the purities $\text{Tr}[\rho^2]$ and $\text{Tr}[\Delta(\rho)^2]$ (notably, the same holds for the tighter second-to-last bound in (9)). The latter two quantities can be measured directly on two copies of the state ρ (assumed unknown), as $\text{Tr}[\rho^{\otimes 2}V]$ and $\text{Tr}[\rho^{\otimes 2}\Delta^{\otimes 2}(V)]$, respectively, with V being the swap operator [30, 48], defined by its action $V|\psi\rangle|\phi\rangle = |\phi\rangle|\psi\rangle$, for all $|\psi\rangle, |\phi\rangle \in \mathbb{C}^d$.

We now show that an *analytical* evaluation of RoC can be obtained for a relevant class of d -dimensional states. Let $\rho \in \mathcal{D}(\mathbb{C}^d)$ be such that there exists an incoherent unitary $U = \sum_j e^{i\phi_j} |j\rangle\langle j|$ such that $(U\rho U^\dagger)_{kl} = |\rho_{kl}|$. One has then $\mathcal{C}_{\mathcal{R}}(\rho) = \mathcal{C}_{\ell_1}(\rho)$ [38], where $\mathcal{C}_{\ell_1}(\rho) = \sum_{k,l} |\rho_{kl}| - 1 = 2 \sum_{k < l} |\rho_{kl}|$ is the ℓ_1 norm of coherence [13]. The class of states for which this equality holds includes, for instance, all one-qubit states ($d = 2$, see Fig. 1), all d -dimensional states with X-shaped density matrix [49–51] (which contain in particular Bell diagonal states of two qubits [31, 52]), and all pure states $|\psi\rangle \in \mathbb{C}^d$. For the latter, writing in general $|\psi\rangle = \sum_{j=0}^{d-1} \psi_j |j\rangle$, we get explicitly $\mathcal{C}_{\mathcal{R}}(|\psi\rangle\langle\psi|) = \mathcal{C}_{\ell_1}(|\psi\rangle\langle\psi|) = (\sum_j |\psi_j|)^2 - 1$ [13].

In particular, maximally coherent states $|\psi^+\rangle$, characterized by $|\psi_j| = \frac{1}{\sqrt{d}} \forall j = 0, \dots, d-1$, have $\mathcal{C}_{\mathcal{R}}(|\psi^+\rangle\langle\psi^+|) = \mathcal{C}_{\ell_1}(|\psi^+\rangle\langle\psi^+|) = d - 1$, that is the maximum possible value for the RoC of any d -dimensional state. One can show [38] in fact that these are the only states which can reach maximal RoC, which positively settles another requirement recently advocated for bona fide measures of coherence [24].

The equivalence between RoC and ℓ_1 norm of coherence breaks down already in dimension $d = 3$. One can prove however the existence of general upper and lower bounds [38],

$$(d-1)^{-1}\mathcal{C}_{\ell_1}(\rho) \leq \mathcal{C}_{\mathcal{R}}(\rho) \leq \mathcal{C}_{\ell_1}(\rho), \quad \forall \rho \in \mathcal{D}(\mathbb{C}^d). \quad (10)$$

Both bounds can be tight. Examples of states saturating the upper bound have been provided already (for instance, all pure states). A family of states saturating the lower bound is instead given by $\rho_p = (1+p)\mathbb{1}/d - p|\psi^+\rangle\langle\psi^+|$, with $0 \leq p \leq \frac{1}{d-1}$, for which $\mathcal{C}_{\ell_1}(\rho_p) = p(d-1)$ and $\mathcal{C}_{\mathcal{R}}(\rho_p) = p$. Nonetheless, the lower bound becomes looser for large values of \mathcal{C}_{ℓ_1} , and one finds $\mathcal{C}_{\mathcal{R}}(\rho) \rightarrow d-1$ for all ρ such that $\mathcal{C}_{\ell_1}(\rho) \rightarrow d-1$ [38].

We are finally ready to provide a direct operational interpretation for the RoC in a metrology context. Consider the following *phase discrimination* (PD) game. Alice prepares a quantum state $\rho \in \mathcal{D}(\mathbb{C}^d)$, which then enters a black box. The black box encodes a phase on ρ by implementing a unitary $U_\phi = \exp(iN\phi)$, with $N = \sum_{j=0}^{d-1} |j\rangle\langle j|$ and $\phi \in \mathbb{R}$, so that the output state is determined by the action of the unitary channel $\mathcal{U}_\phi(\rho) := U_\phi \rho U_\phi^\dagger$. We can think of N as a Hamiltonian for the system with equispaced spectrum, assuming unit spacing without loss of generality. Suppose one of m phases $\{\phi_k\}_{k=0}^{m-1}$ can be applied, each with a prior probability p_k . Any collection of pairs $\{(p_k, \phi_k)\}_{k=0}^{m-1} =: \Theta$ defines a PD game, where Alice's goal is that of guessing correctly the phase that was actually imprinted on the state. To this end, she performs a generalized measurement with elements $\{M_k\}$ (satisfying $M_k \geq 0$, $\sum_k M_k = \mathbb{1}$) on the output state $\mathcal{U}_\phi(\rho)$ after the black box. Optimizing over all measurements, the maximal probability of success depends on the game Θ and the input state ρ , and is given by $p_\Theta^{\text{succ}}(\rho) = \max_{\{M_k\}} \sum_k p_k \text{Tr}[U_{\phi_k} \rho U_{\phi_k}^\dagger M_k]$.

Suppose now Alice's input state is incoherent, $\rho \equiv \delta \in \mathcal{I}$. Since every unitary channel \mathcal{U}_ϕ leaves any such state invariant, $\mathcal{U}_\phi(\delta) = \delta$, the best strategy for Alice is always to cast the guess k^{max} corresponding to the phase with the highest prior probability $p_{k^{\text{max}}} := \max_k p_k$. This results in an optimal probability of success for any incoherent state given by $p_\Theta^{\text{succ}}(\mathcal{I}) := p_{k^{\text{max}}}$, which can be achieved even without actually probing the channel, just by a fixed guess.

It is clear that, by preparing a coherent state $\rho \notin \mathcal{I}$, Alice can expect to do better. What is less obvious yet more remarkable, is that the maximum advantage achievable by using ρ as opposed to any incoherent probe δ , in all possible PD games, is quantified exactly by the RoC of ρ . More precisely [38]:

$$\max_{\Theta} \frac{p_\Theta^{\text{succ}}(\rho)}{p_\Theta^{\text{succ}}(\mathcal{I})} = 1 + \mathcal{C}_{\mathcal{R}}(\rho). \quad (11)$$

The maximum is achieved for the PD game $\Theta^* \equiv \{(\frac{1}{d}, \frac{2\pi k}{d})\}_{k=0}^{d-1}$. Therefore $\mathcal{C}_{\mathcal{R}}(\rho)$ exactly quantifies, in particular, how useful the state ρ is for reliable decoding and transmission of messages encoded by generalized phase channels $\rho \mapsto Z^k \rho Z^{\dagger k}$, with $Z|j\rangle = \exp(i\frac{2\pi}{d}j)|j\rangle$. These channels feature in several quantum information tasks such as quantum error correction [53], cloning [54], and dense coding [55, 56]. This suggests a prominent role of coherence, specifically measured by the RoC, in quantum communication.

We notice that one can consider more general channel discrimination (CD) games, where each game is associated with a set of pairs $\{(p_k, \Lambda_k)\}_{k=0}^{m-1} =: \Upsilon$, with $\{\Lambda_k\}$ a set of m (generally nonunitary) channels. For each CD game Υ , Alice's

goal is still that of discriminating which Λ_k gets applied by a black box to an input ρ , and she succeeds with optimal probability $p_\Upsilon^{\text{succ}}(\rho) = \max_{\{M_k\}} \sum_k p_k \text{Tr}[\Lambda_k(\rho)M_k]$, where we optimize over measurements similarly as before. We observe that the definition of RoC, and in particular the existence of the decomposition (2), is such that for any CD game Υ it holds $p_\Upsilon^{\text{succ}}(\rho) \leq (1 + \mathcal{C}_{\mathcal{R}}(\rho))p_\Upsilon^{\text{succ}}(\mathcal{I})$, where $p_\Upsilon^{\text{succ}}(\mathcal{I})$ is the best probability of success by using as input an arbitrary incoherent state. Notice that in general, $p_\Upsilon^{\text{succ}}(\mathcal{I})$ can be higher than $p_{k^{\text{max}}}$, because the channels Λ_k may act nontrivially on incoherent states. Nonetheless, if one focuses on a subclass of CD games $\{\Upsilon^*\}$ containing the PD game Θ^* , i.e., $\Theta^* \in \{\Upsilon^*\}$, one gets: $\max_{\Upsilon \in \{\Upsilon^*\}} \frac{p_\Upsilon^{\text{succ}}(\rho)}{p_\Upsilon^{\text{succ}}(\mathcal{I})} = \frac{p_{\Theta^*}^{\text{succ}}(\rho)}{p_{\Theta^*}^{\text{succ}}(\mathcal{I})} = 1 + \mathcal{C}_{\mathcal{R}}(\rho)$. The RoC $\mathcal{C}_{\mathcal{R}}(\rho)$ thus quantifies the maximum achievable advantage in any CD task in which the phase channels Z^k are some of the possible channels applied to a probe ρ by a black box.

It will be a worthy development to extend this analysis to the scenario of *assisted* CD games, where the collaboration of a correlated party Bob may further increase Alice's probability of success in the discrimination. A similar setting has been studied recently for the task of distilling coherence [57]. It will be then especially interesting to investigate the interplay between Alice's local (assisted) RoC and the robustness of entanglement [32, 33] in the state shared between Alice and Bob. Notice that, in different channel discrimination problems, the resource roles of the robustness of entanglement [58] and steering [34] have been acknowledged.

We conclude by announcing once again that the definition (1) can be extended to a more abstract notion of *robustness of asymmetry*, in which the free states are identified as those invariant with respect to the action of a group [18]. Then, suitable adaptations of all the properties demonstrated above in Eqs. (3)–(9) carry over [38], including the operational interpretation in the context of channel discrimination games. The approach pursued in this Letter, based on the generalized notion of robustness, appears accordingly quite versatile to define and validate insightful quantifiers of resources in quantum physics [27] and beyond [26, 59], as demonstrated here for the fundamental case of quantum coherence.

Acknowledgments. This work has received funding from the European Research Council (ERC StG GQCOP, Grant No. 637352), from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie (Grant Agreement No. 661338), and the Erasmus+ Programme. We thank D. Girolami, I. Marvian, M. B. Plenio, and A. Streltsov for useful discussions.

-
- [1] B. Hensen, H. Bernien, A. Dréau, A. Reiserer, N. Kalb, M. Blok, J. Ruitenberg, R. Vermeulen, R. Schouten, C. Abellán, W. Amaya, V. Pruneri, M. W. Mitchell, M. Markham, D. Twitchen, D. Elkouss, S. Wehner, T. Tamini, and R. Hanson, *Nature* **526**, 682 (2015).
 - [2] M. Giustina, M. A. M. Versteegh, S. Wengerowsky, J. Hand-

- steiner, A. Hochrainer, K. Phelan, F. Steinlechner, J. Kofler, J.-A. Larsson, C. Abellán, W. Amaya, V. Pruneri, M. W. Mitchell, J. Beyer, T. Gerrits, A. E. Lita, L. K. Shalm, S. W. Nam, T. Scheidl, R. Ursin, B. Wittmann, and A. Zeilinger, *Phys. Rev. Lett.* **115**, 250401 (2015).
- [3] L. K. Shalm, E. Meyer-Scott, B. G. Christensen, P. Bierhorst, M. A. Wayne, M. J. Stevens, T. Gerrits, S. Glancy, D. R. Hamel, M. S. Allman, K. J. Coakley, S. D. Dyer, C. Hodge, A. E. Lita, V. B. Verma, C. Lambrocco, E. Tortorici, A. L. Migdall, Y. Zhang, D. R. Kumor, W. H. Farr, F. Marsili, M. D. Shaw, J. A. Stern, C. Abellán, W. Amaya, V. Pruneri, T. Jennewein, M. W. Mitchell, P. G. Kwiat, J. C. Bienfang, R. P. Mirin, E. Knill, and S. W. Nam, *Phys. Rev. Lett.* **115**, 250402 (2015).
- [4] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information* (Cambridge university press, 2010).
- [5] *A little bit, better* (The Economist, 20th June 2015).
- [6] J. P. Dowling and G. J. Milburn, *Phil. Trans. Roy. Soc. A* **361**, 1655 (2003).
- [7] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [8] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, *Rev. Mod. Phys.* **84**, 1655 (2012).
- [9] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, *Rev. Mod. Phys.* **86**, 419 (2014).
- [10] S. Lloyd, *J. Phys.: Conf. Ser.* **302**, 012037 (2011).
- [11] S. F. Huelga and M. B. Plenio, *Contemp. Phys.* **54**, 181 (2013).
- [12] J. Åberg, “Quantifying superposition,” (2006), arXiv:quant-ph/0612146.
- [13] T. Baumgratz, M. Cramer, and M. B. Plenio, *Phys. Rev. Lett.* **113**, 140401 (2014).
- [14] F. Grosshans, G. Van Assche, J. Wenger, R. Brouri, N. J. Cerf, and P. Grangier, *Nature* **421**, 238 (2003).
- [15] V. Giovannetti, S. Lloyd, and L. Maccone, *Nat. Photon.* **5**, 222 (2011).
- [16] M. Lostaglio, D. Jennings, and T. Rudolph, *Nature Commun.* **6**, 6383 (2015).
- [17] V. Narasimhachar and G. Gour, *Nature Commun.* **6**, 7689 (2015).
- [18] I. Marvian and R. W. Spekkens, *New J. Phys.* **15**, 033001 (2013).
- [19] F. Levi and F. Mintert, *New J. Phys.* **16**, 033007 (2014).
- [20] I. Marvian, R. W. Spekkens, and P. Zanardi, (2015), arXiv:1510.06474.
- [21] A. Streltsov, U. Singh, H. S. Dhar, M. N. Bera, and G. Adesso, *Phys. Rev. Lett.* **115**, 020403 (2015).
- [22] S. Du, Z. Bai, and Y. Guo, *Phys. Rev. A* **91**, 052120 (2015).
- [23] A. Winter and D. Yang, (2015), arXiv:1506.07975.
- [24] Y. Peng, Y. Jiang, and H. Fan, (2015), arXiv:1511.02576.
- [25] A. Streltsov, “Genuine quantum coherence,” (2015), arXiv:1511.08346.
- [26] B. Coecke, T. Fritz, and R. Spekkens, “A mathematical theory of resources,” (2014), arXiv:1409.5531.
- [27] F. G. S. L. Brandão and G. Gour, *Phys. Rev. Lett.* **115**, 070503 (2015).
- [28] B. Yadin, J. Ma, D. Girolami, M. Gu, and V. Vedral, “Quantum processes which do not use coherence,” (2015), arXiv:1512.02085.
- [29] I. Marvian and R. W. Spekkens, *Nat. Commun.* **5**, 3821 (2014).
- [30] D. Girolami, *Phys. Rev. Lett.* **113**, 170401 (2014).
- [31] T. R. Bromley, M. Cianciaruso, and G. Adesso, *Phys. Rev. Lett.* **114**, 210401 (2015).
- [32] G. Vidal and R. Tarrach, *Phys. Rev. A* **59**, 141 (1999).
- [33] M. Steiner, *Phys. Rev. A* **67**, 054305 (2003).
- [34] M. Piani and J. Watrous, *Phys. Rev. Lett.* **114**, 060404 (2015).
- [35] J. Geller and M. Piani, *J. Phys. A: Math. Theor.* **47** (2014).
- [36] The Supplemental Material contains Matlab code to evaluate the RoC for arbitrary quantum states, as well as from experimental data.
- [37] C.-M. Li, N. Lambert, Y.-N. Chen, G.-Y. Chen, and F. Nori, *Sci. Rep.* **2**, 885 (2012).
- [38] M. Piani, M. Cianciaruso, T. R. Bromley, C. Napoli, N. Johnston, and G. Adesso, “Robustness of asymmetry and coherence of quantum states”, coordinated submission to *Phys. Rev. A* (2016).
- [39] J. Eisert, F. G. Brandão, and K. M. Audenaert, *New J. Phys.* **9**, 46 (2007).
- [40] O. Gühne and G. Tóth, *Physics Reports* **474**, 1 (2009).
- [41] Any W such that $\Delta(W) = 0$ satisfies $\text{Tr}[\delta W] = \text{Tr}[\delta \Delta(W)] = 0$ for any incoherent state δ , so that, if $\text{Tr}[\rho W] \neq 0$, then ρ must have some coherence. Nonetheless, meeting the exact equality condition $\Delta(W) = 0$ is experimentally challenging, while the condition $\Delta(W) \geq 0$ is experimentally friendlier, and sets the basis for a practical verification of coherence.
- [42] L. Vandenberghe and S. Boyd, *SIAM review* **38**, 49 (1996).
- [43] F. G. S. L. Brandão, *Phys. Rev. A* **72**, 022310 (2005).
- [44] MATLAB, *version 8.5.0 (R2015a)* (The MathWorks Inc., Natick, Massachusetts, 2015).
- [45] M. Grant and S. Boyd, “CVX: Matlab software for disciplined convex programming, version 2.1,” <http://cvxr.com/cvx> (2014).
- [46] M. Grant and S. Boyd, in *Recent Advances in Learning and Control*, Lecture Notes in Control and Information Sciences, edited by V. Blondel, S. Boyd, and H. Kimura (Springer-Verlag Limited, 2008) pp. 95–110, http://stanford.edu/~boyd/graph_dcp.html.
- [47] M. Sarovar, A. Ishizaki, G. R. Fleming, and K. B. Whaley, *Nat. Phys.* **6**, 462 (2010).
- [48] A. K. Ekert, C. M. Alves, D. K. Oi, M. Horodecki, P. Horodecki, and L. C. Kwek, *Physical review letters* **88**, 217901 (2002).
- [49] S. Vinjanampathy and A. R. P. Rau, *Phys. Rev. A* **82**, 032336 (2010).
- [50] S. M. Hashemi Rafsanjani, M. Huber, C. J. Broadbent, and J. H. Eberly, *Phys. Rev. A* **86**, 062303 (2012).
- [51] Generalized X-states can be defined as $\rho = \sum_{k=0}^{d/2} \rho_k$ if d is even, and $\rho = \sum_{k=0}^{\lfloor d/2 \rfloor} \rho_k + \rho_c$ if d is odd, with $\rho_k = \rho_{kk} |k\rangle\langle k| + \rho_{k,d-1-k} |k\rangle\langle d-1-k| + \rho_{d-1-k,k} |d-1-k\rangle\langle k| + \rho_{d-1-k,d-1-k} |d-1-k\rangle\langle d-1-k|$, and $\rho_c = \rho_{\lfloor d/2 \rfloor + 1, \lfloor d/2 \rfloor + 1} |\lfloor d/2 \rfloor + 1\rangle\langle \lfloor d/2 \rfloor + 1|$.
- [52] R. Horodecki and M. Horodecki, *Phys. Rev. A* **54**, 1838 (1996).
- [53] D. Gottesman, *Chaos, Solitons & Fractals* **10**, 1749 (1999).
- [54] N. J. Cerf, *J. Mod. Opt.* **47**, 197 (2000).
- [55] T. Hiroshima, *J. Phys. A: Math. Gen.* **34**, 6907 (2001).
- [56] R. F. Werner, *J. Phys. A: Math. Gen.* **34**, 7094 (2001).
- [57] E. Chitambar, A. Streltsov, S. Rana, M. N. Bera, G. Adesso, and M. Lewenstein, (2015), arXiv:1507.08171v1.
- [58] M. Piani and J. Watrous, *Phys. Rev. Lett.* **102**, 250501 (2009).
- [59] L. del Rio, L. Krämer, and R. Renner, “Resource theories of knowledge,” (2015), arXiv:1511.08818.